
### Class 07: Outline

- Hour 1:
- Conductors & Insulators Expt. 2: Electrostatic Force Hour 2:
  - Capacitors

Last Time: Gauss's Law

### **Gauss's Law**



$$\Phi_E = \bigoplus_{\substack{\text{closed}\\\text{surface S}}} \vec{\mathbf{E}} \cdot d\vec{\mathbf{A}} = \frac{Q_{enc}}{\mathcal{E}_0}$$

In practice, use symmetry:

- Spherical (r)
- Cylindrical (r, l)
- Planar (Pillbox, A)

# Conductors

### **Conductors and Insulators**


A conductor contains charges that are free to move (electrons are weakly bound to atoms) Example: metals

An insulator contains charges that are NOT free to move (electrons are strongly bound to atoms) Examples: plastic, paper, wood

### Conductors


Conductors have free charges

- $\rightarrow$  E must be zero inside the conductor
- → Conductors are equipotential objects




### Equipotentials

### **Topographic Maps**



### **Equipotential Curves**





All points on equipotential curve are at same potential. Each curve represented by V(x,y) = constant


# PRS Question: Walking down a mountain

# **Direction of Electric Field E**

#### E is perpendicular to all equipotentials







**Constant E field** 

**Point Charge** 

**Electric dipole** 


# **Properties of Equipotentials**

- E field lines point from high to low potential
- E field lines perpendicular to equipotentials
  - Have no component along equipotential
  - No work to move along equipotential

### **Conductors in Equilibrium**


- Conductors are equipotential objects:
- 1) E = 0 inside
- 2) Net charge inside is 0
- 3) E perpendicular to surface
- 4) Excess charge on surface

$$E = \sigma / \varepsilon_0$$



### **Conductors in Equilibrium**

Put a net positive charge anywhere inside a conductor, and it will move to the surface to get as far away as possible from the other charges of like sign.




http://ocw.mit.edu/ans7870/8/8.02T/f04/visualizations/electrostatics/34-pentagon/34pentagon320.html

### **Expt. 2: Electrostatic Force**



### **Expt. 2: Electrostatic Force**




http://ocw.mit.edu/ans7870/8/8.02T/f04/visualizations/electrostatics/36-electrostaticforce/36-esforce320.html


### **Experiment 2: Electrostatic Force**

### **Capacitors and Capacitance**

### **Capacitors: Store Electric Energy**

Capacitor: two isolated conductors with equal and opposite charges Q and potential difference  $\Delta V$  between them.






Units: Coulombs/Volt or Farads

# **Parallel Plate Capacitor** E = 0+O $=\sigma A$ EE = 0


### **Parallel Plate Capacitor**

When you put opposite charges on plates, charges move to the inner surfaces of the plates to get as close as possible to charges of the opposite sign




http://ocw.mit.edu/ans7870/8/8.02T/f04/visuali zations/electrostatics/35-capacitor/35capacitor320.html

# Calculating E (Gauss's Law)




Note: We only "consider" a single sheet! Doesn't the other sheet matter?

### **Alternate Calculation Method**



$$E = \frac{\sigma}{2\varepsilon_0} + \frac{\sigma}{2\varepsilon_0} = \frac{\sigma}{\varepsilon_0} = \frac{Q}{A\varepsilon_0}$$

### **Parallel Plate Capacitor**




C depends only on geometric factors A and d P07 -24

bottom

Demonstration: Big Capacitor

### **Spherical Capacitor**

Two concentric spherical shells of radii a and b



What is E?

Gauss's Law  $\rightarrow$  E  $\neq$  0 only for *a* < *r* < *b*, where it looks like a point charge:

$$\vec{\mathbf{E}} = \frac{Q}{4\pi\varepsilon_0 r^2} \hat{\mathbf{r}}$$

### **Spherical Capacitor**

$$\Delta V = -\int_{inside}^{outside} \vec{\mathbf{E}} \cdot d\vec{\mathbf{S}} = -\int_{a}^{b} \frac{Q\hat{\mathbf{r}}}{4\pi\varepsilon_{0}r^{2}} \cdot dr\,\hat{\mathbf{r}} = \frac{Q}{4\pi\varepsilon_{0}} \left(\frac{1}{b} - \frac{1}{a}\right)$$

Is this positive or negative? Why?

$$C = \frac{Q}{|\Delta V|} = \frac{4\pi\varepsilon_0}{\left(a^{-1} - b^{-1}\right)}$$

$$-Q$$
  
+ $Q/a$   
b

For an isolated spherical conductor of radius *a*:  $C = 4\pi \varepsilon_0 a$ 

### **Capacitance of Earth**

For an isolated spherical conductor of radius a:

$$C = 4\pi\varepsilon_0 a$$

$$\varepsilon_0 = 8.85 \times 10^{-12} \text{ F/m}$$
  $a = 6.4 \times 10^6 \text{ m}$ 

$$C = 7 \times 10^{-4} \,\mathrm{F} = 0.7 \,\mathrm{mF}$$

A Farad is REALLY BIG! We usually use pF (10<sup>-12</sup>) or nF (10<sup>-9</sup>)

### **1 Farad Capacitor**


How much charge?  $Q = C |\Delta V|$  = (1F)(12V) = 12C

# PRS Question: Changing C Dimensions

# Demonstration: Changing C Dimensions

# **Energy Stored in Capacitor**


# **Energy To Charge Capacitor**



- 1. Capacitor starts uncharged.
- 2. Carry +dq from bottom to top. Now top has charge q = +dq, bottom -dq
- 3. Repeat
- 4. Finish when top has charge q = +Q, bottom -Q

# **Work Done Charging Capacitor**

At some point top plate has +q, bottom has -qPotential difference is  $\Delta V = q / C$ Work done lifting another dq is  $dW = dq \Delta V$ 




# **Work Done Charging Capacitor**

So work done to move dq is:

$$dW = dq \,\Delta V = dq \,\frac{q}{C} = \frac{1}{C} q \,dq$$

Total energy to charge to q = Q:



**Energy Stored in Capacitor** 

Since 
$$C = \frac{Q}{\left|\Delta V\right|}$$

$$U = \frac{Q^2}{2C} = \frac{1}{2}Q|\Delta V| = \frac{1}{2}C|\Delta V|^2$$

Where is the energy stored???

**Energy Stored in Capacitor** Energy stored in the E field! **Parallel-plate capacitor:**  $C = \frac{\varepsilon_o A}{d}$  and V = Ed $U = \frac{1}{2}CV^2 = \frac{1}{2}\frac{\varepsilon_o A}{d} (Ed)^2 = \frac{\varepsilon_o E^2}{2} \times (Ad) = u_E \times (volume)$ 

$$u_E = E$$
 field energy density  $= \frac{\varepsilon_o E^2}{2}$ 

### **1 Farad Capacitor - Energy**

How much energy?

$$U = \frac{1}{2} C |\Delta V|^{2}$$
  
=  $\frac{1}{2} (1 F) (1 2 V)^{2}$   
= 72 J

Compare to capacitor charged to 3kV:

$$U = \frac{1}{2} C \left| \Delta V \right|^2 = \frac{1}{2} (100 \,\mu\,\mathrm{F}) (3\,\mathrm{k\,V})^2$$
$$= \frac{1}{2} (1 \times 10^{-4}\,\mathrm{F}) (3 \times 10^3\,\mathrm{V})^2 = 450\,\mathrm{J}$$

PRS Question: Changing C Dimensions Energy Stored

## Demonstration: Dissectible Capacitor